arXiv:1406.1829v2 [math.GR] 24 May 2016

HAUSDORFF DIMENSION IN R-ANALYTIC PROFINITE GROUPS

GUSTAVO A. FERNÁNDEZ-ALCOBER, EUGENIO GIANNELLI, AND JON GONZÁLEZ-SÁNCHEZ

Abstract. We study the Hausdorff dimension of R-analytic subgroups in an R-analytic profinite group, where R is a pro-p ring whose associated graded ring is an integral domain. In particular, we prove that the set of such Hausdorff dimensions is a finite subset of the rational numbers.

1. Introduction

The study of Hausdorff dimension in profinite groups was initiated by Abercrombie in [1], and has attracted special attention in recent times; see, for example, [2, 3, 4, 7, 9, 11]. If G is a countably based infinite profinite group, a *(normal) filtration* of G is a descending series $\{G_n\}_{n\in\mathbb{N}}$ of (normal) open subgroups of G which is a base of neighbourhoods of 1. We can define a metric on G by

$$d(x,y) = \inf\{|G:G_n|^{-1} \mid xy^{-1} \in G_n\},$$

and the topology induced by d coincides with the original topology in G. The metric d defines a Hausdorff dimension function on all subsets of G, which we denote by $\mathrm{hDim}_{\{G_n\}}$. As shown in [4, Theorem 2.4], if the filtration $\{G_n\}_{n\in\mathbb{N}}$ is normal and H is a closed subgroup of G, then

$$\mathrm{hDim}_{\{G_n\}}(H) = \liminf_{n\to\infty} \frac{\log|H : H\cap G_n|}{\log|G:G_n|}.$$

In [4], Barnea and Shalev studied the Hausdorff dimension in p-adic analytic pro-p groups with respect to the filtration $\{G^{p^n}\}_{n\in\mathbb{N}}$, and proved that

$$\mathrm{hDim}_{\{G^{p^n}\}}(H) = \frac{\dim(H)}{\dim(G)}$$

for every closed subgroup H of G. Here, if G is an analytic group, $\dim(G)$ denotes the dimension of G as an analytic manifold.

In this paper we study Hausdorff dimension in R-analytic profinite groups, where R is a pro-p ring whose associated graded ring is an integral domain. This is the natural setting for working with analytic groups over pro-p rings, see [5, Chapter 13]. If G is an R-analytic profinite group, then G is virtually

2010 *Mathematics Subject Classification.* Primary 20E18; Secondary 28A78.

Key words and phrases. Profinite groups, analytic groups, Hausdorff dimension.

The first and third authors are supported by the Spanish Government, grants MTM2011-28229-C02 and MTM2014-53810-C2-2-P, and by the Basque Government, grants IT753-13 and IT974-16. The second author gratefully acknowledges financial support by the GRECA research group and by the ERC Advanced Grant 291512.

a pro-p group, but the subgroups G^{p^n} need not to form a filtration, since they need not to be open in G. We consider instead the natural filtration induced by a standard open subgroup of G. As we will see, the Hausdorff dimension of a closed subgroup H of G is the same with respect to the natural filtrations induced by all open standard subgroups of G. This allows us to define the concept of *standard Hausdorff dimension* of H, denoted by $\mathrm{hDim}_{\mathrm{Std}}(H)$. We get the following theorem, in the spirit of the result of Barnea and Shalev.

Main Theorem. *Let G be an R-analytic profinite group and let H be an R-analytic subgroup of G. Then*

$$\mathrm{hDim}_{\mathrm{Std}}(H) = \frac{\dim(H)}{\dim(G)}. \tag{1}$$

An R-analytic subgroup H of G is a subgroup that is also an R-analytic submanifold of G; then H is closed in G. A crucial remark is that while the converse is true in a p-adic analytic group, i.e. every closed subgroup is analytic, it need not hold for an arbitrary pro-p ring R. For example, $\mathbb{F}_p[[t^d]]$ is a closed subgroup of $\mathbb{F}_p[[t]]$ for every positive integer d, but it is not analytic for $d > 1$, even if it is an $\mathbb{F}_p[[t]]$-analytic group in its own right.

Observe that (1) implies that the *R-analytic spectrum* of G, defined by

$$\mathrm{Spec}_R(G) = \{\mathrm{hDim}_{\mathrm{Std}}(H) \mid H \text{ is an } R\text{-analytic subgroup of } G\},$$

is finite and consists of rational values. On the other hand, the spectrum of $\mathbb{F}_p[[t]]$ corresponding to all closed subgroups is the full interval $[0, 1]$, as shown in [4, Lemma 4.1]. Thus our main theorem is pointing to the fact that most closed subgroups of $\mathbb{F}_p[[t]]$ are non-analytic.

Notation. We write $X^{(d)}$ to denote the cartesian product of d copies of a set X. The symbol $\subseteq_o$ indicates that a subset of a topological space is open.

2. Preliminaries

Throughout this paper R is a pro-p ring whose associated graded ring is an integral domain. Hence R is an integral domain, and we write K for its field of fractions. If $\mathfrak{m}$ is the maximal ideal of R, then $R/\mathfrak{m}$ is a finite field of characteristic p. Set $H(n) = \dim_{R/\mathfrak{m}}(\mathfrak{m}^n/\mathfrak{m}^{n+1})$ for all $n \in \mathbb{N}$. For large enough n, $H(n)$ coincides with a polynomial $P(n)$, called the *Hilbert polynomial* of R [6, Chapter 8, Theorem C].

Let G be an R-analytic group. Without loss of generality, we assume that the manifold structure is given by a full atlas. A subgroup H of G is an *analytic subgroup* if it is also an analytic submanifold of G. We adopt Serre's definition of submanifold [10, Section 3.11]. Every analytic subgroup is closed in G. The converse is not true in general, but every open subgroup H of G (actually, any open subset) is a submanifold when considered with the restrictions of the charts of G, and $\dim H = \dim G$.

An R-analytic group S with a global chart given by a homeomorphism $\phi : S \to (\mathfrak{m}^N)^{(d)}$ is called an *R-standard group* of level N if $\phi(1) = 0$ and, for all $j \in \{1, \dots, d\}$, there exist $F_j(X, Y) \in R[[X, Y]]$ (where X and Y are

d-tuples of indeterminates), without constant term, such that

$$\phi(xy) = F(\phi(x), \phi(y)), \quad \text{for every } x, y \in S. \tag{2}$$

Here, $F = (F_1, \dots, F_d)$ is called the *formal group law* associated to S. In these circumstances, we also say that (S, ϕ) is a standard group.

By using (2), F defines a new group structure on $(\mathfrak{m}^N)^{(d)}$, other than its natural additive structure, and then ϕ is an isomorphism between S and $((\mathfrak{m}^N)^{(d)}, F)$. Every R-analytic group contains an open (and so analytic) R-standard subgroup S [5, Theorem 13.20]. As a consequence, profinite R-analytic groups are countably based.

The formal group law of a standard group satisfies the following result [5, pp. 331-334].

Lemma 2.1. *Let F be a formal group law of dimension d associated to a standard group S. Then*

$$F(X, Y) = X + Y + G(X, Y),$$

where every monomial involved in G has total degree at least 2, and contains a non-zero power of X_r and of Y_s for some $r, s \in \{1, \dots, d\}$. Moreover,

$$F(X, I(X)) = 0 = F(I(X), X)$$

for some $I(X) = -X + H(X) \in R[[X]]$, and every monomial involved in H has total degree at least 2.

It follows that $(\mathfrak{m}^{N+n})^{(d)}$ is a subgroup of $((\mathfrak{m}^N)^{(d)}, F)$ for all $n \geq 0$. This allows us to introduce a special type of filtrations in an R-analytic group.

Definition 2.2. Let G be an R-analytic group and let S be a standard open subgroup of G, with global chart (S, ϕ). Then for every positive integer $n \geq 0$, and for every subset $A \subseteq S$, we define

$$\mathfrak{m}_\phi^n A = \phi^{-1}(\mathfrak{m}^n \phi(A)).$$

We say that $\{\mathfrak{m}_\phi^n S\}_{n \in \mathbb{N}}$ is the *natural filtration* of G induced by S.

Observe that

$$\mathfrak{m}_\phi^n S = \phi^{-1}((\mathfrak{m}^{N+n})^{(d)}),$$

which implies that $\mathfrak{m}_\phi^n S$ is an open subgroup of G, and also that $\{\mathfrak{m}_\phi^n S\}_{n \in \mathbb{N}}$ is a filtration of G. Actually, we have $\mathfrak{m}_\phi^n S \trianglelefteq S$ (see [5, Proposition 13.22]).

Lemma 2.3. *Let (S, ϕ) be an R-standard group. Then:*

(i) *For all $x, y \in S$ and $n \geq N$, we have $\phi(xy^{-1}) \in (\mathfrak{m}^n)^{(d)}$ if and only if $\phi(x) - \phi(y) \in (\mathfrak{m}^n)^{(d)}$.*
(ii) *For every $n \geq 0$, we have $|S : \mathfrak{m}_\phi^n S| = |(\mathfrak{m}^N)^{(d)} : (\mathfrak{m}^{N+n})^{(d)}| = q^{df(n)}$, where $f(n) = \sum_{i=N}^{N+n-1} H(i)$.*
(iii) *ϕ is an isometry between the group S with the metric induced by the filtration $\{\mathfrak{m}_\phi^n S\}_{n \in \mathbb{N}}$ and the group $((\mathfrak{m}^N)^{(d)}, +)$ with the metric induced by the filtration $\{(\mathfrak{m}^{N+n})^{(d)}\}_{n \in \mathbb{N}}$.*

Proof. (i) Since $\phi(1) = 0$, we may assume that $x \neq y$. Let $k \in \mathbb{N}$ be such that $\phi(xy^{-1}) \in (\mathfrak{m}^k)^{(d)} \setminus (\mathfrak{m}^{k+1})^{(d)}$. Then since

$$\phi(x) = \phi(xy^{-1}y) = F(\phi(xy^{-1}), \phi(y)) = \phi(xy^{-1}) + \phi(y) + G(\phi(xy^{-1}), \phi(y)),$$

by Lemma 2.1, we have

$$\phi(x) - \phi(y) \equiv \phi(xy^{-1}) \pmod{(\mathfrak{m}^{k+1})^{(d)}}.$$

Hence also $\phi(x) - \phi(y) \in (\mathfrak{m}^k)^{(d)} \smallsetminus (\mathfrak{m}^{k+1})^{(d)}$, and (i) follows.

(ii) Observe that (i) implies that

$$xy^{-1} \in \mathfrak{m}_\phi^n S \iff \phi(x) - \phi(y) \in (\mathfrak{m}^{N+n})^{(d)}, \tag{3}$$

or what is the same,

$$x \cdot \mathfrak{m}_\phi^n S = y \cdot \mathfrak{m}_\phi^n S \iff \phi(x) + (\mathfrak{m}^{N+n})^{(d)} = \phi(y) + (\mathfrak{m}^{N+n})^{(d)}.$$

This proves (ii), since ϕ is a bijection.

(iii) According to the definition of the metric associated to a fitration, it is clear that (ii) and (3) together imply that ϕ is an isometry. □

Hausdorff dimension can be defined for any subset of a metric space, see [8, Chapter 2] for its definition and main properties. We need the following two lemmas about Hausdorff dimension in countably based profinite groups.

Lemma 2.4. *Let G be a countably based profinite group with filtration $\{G_n\}_{n\in\mathbb{N}}$. Let H be a closed subgroup of G, and let U be a non-empty open subset of H. Then*

$$\mathrm{hDim}_{\{G_n\}}(H) = \mathrm{hDim}_{\{G_n\}}(U).$$

Proof. The proof is a straightforward consequence of [8, Section 2.2]. □

Let G be a countably based profinite group and let S be an open subgroup of G. If $\{S_n\}_{n\in\mathbb{N}}$ is a filtration of S, we can calculate the Hausdorff dimension of $X \subseteq S$ with respect to the metric induced by $\{S_n\}_{n\in\mathbb{N}}$ in S or in G, which we denote by $\mathrm{hDim}^S_{\{S_n\}}(X)$ and $\mathrm{hDim}^G_{\{S_n\}}(X)$, respectively. Our next lemma shows that there is no need to make this distinction in the notation.

Lemma 2.5. *Let G be a countably based profinite group and let S be an open subgroup of G. If $\{S_n\}_{n\in\mathbb{N}}$ is a filtration of S, then*

$$\mathrm{hDim}^G_{\{S_n\}}(X) = \mathrm{hDim}^S_{\{S_n\}}(X)$$

for every $X \subseteq S$.

Proof. Let d_S and d_G be the metrics induced by $\{S_n\}_{n\in\mathbb{N}}$ in S and G. Then $d_S(x, y) = |G : S|\, d_G(x, y)$ for all $x, y \in S$, and the identity map from (S, d_S) to (S, d_G) is bi-Lipschitz. Now the result follows from [8, Corollary 2.4]. □

3. Proof of the main theorem

In this section we first prove that the Hausdorff dimension of a closed subgroup in an analytic profinite group with respect to a natural filtration is independent of the standard subgroup. Then we prove the main theorem of our paper about Hausdorff dimension of analytic subgroups.

Theorem 3.1. *Let G be an R-analytic profinite group, and let (S, ϕ) and (T, ψ) be two open standard subgroups of G. Then*

$$\mathrm{hDim}_{\{\mathfrak{m}_\phi^n S\}}(H) = \mathrm{hDim}_{\{\mathfrak{m}_\psi^n T\}}(H)$$

for every closed subgroup H of G.

Proof. Let us write S_n and T_n for $\mathfrak{m}_\phi^n S$ and $\mathfrak{m}_\psi^n T$, and $N(S)$ and $N(T)$ for the levels of S and T. We first show that there exist non-negative integers a and b such that

$$S_{n+a} \leq T_n \leq S_{n-b} \tag{4}$$

for every $n \geq b$. Since the charts ϕ and ψ belong to the full atlas of G, the two functions $\psi \circ \phi^{-1}|_{\phi(S\cap T)}$ and $\phi \circ \psi^{-1}|_{\psi(S\cap T)}$ are analytic. Since $\phi(S \cap T)$ is open in $\phi(S)$, it follows that $\psi \circ \phi^{-1}$ can be evaluated in $(\mathfrak{m}^\ell)^{(d)}$ for some ℓ. By [5, Lemma 6.45], and observing that $\psi \circ \phi^{-1}(0) = 0$, there exists a natural number k such that

$$\psi \circ \phi^{-1}((\mathfrak{m}^{n+k})^{(d)}) \subseteq (\mathfrak{m}^n)^{(d)}$$

for every $n \geq 0$. This implies that $S_{n+a} \leq T_n$ for $n \geq 0$, by choosing $a = \max\{k - N(S) + N(T), 0\}$. Arguing similarly with $\phi \circ \psi^{-1}$, we get (4).

Now, by [4, Theorem 2.4], we have

$$\begin{aligned}
\mathrm{hDim}_{\{T_n\}}(H) &= \liminf_{n\to\infty} \frac{\log|H : H \cap T_n|}{\log|G : T_n|} \\
&\leq \liminf_{n\to\infty} \frac{\log|H : H \cap S_{n+a}|}{\log|G : S_{n+a}| - \log|T_n : S_{n+a}|} \\
&\leq \liminf_{n\to\infty} \frac{\log|H : H \cap S_{n+a}|}{\log|G : S_{n+a}|} \cdot \left(1 - \frac{\log|S_{n-b} : S_{n+a}|}{\log|G : S_{n+a}|}\right)^{-1} \\
&= \mathrm{hDim}_{\{S_n\}}(H).
\end{aligned}$$

The last equality holds because, by (ii) of Lemma 2.3, for n large enough we have

$$\log_q|S_{n-b} : S_{n+a}| = \sum_{i=N+n-b}^{N+n+a-1} P(i),$$

which is polynomial in n of degree $\deg P$; on the other hand, $\log_q|G : S_{n+a}|$ is asymptotically equivalent to

$$\log_q|G : S| + \sum_{i=N}^{N+n+a-1} P(i)$$

and, by the Euler-MacLaurin formula, this sum is a polynomial in n of degree $\deg P + 1$. By swapping S and T, the result follows. □

Definition 3.2. Let G be an R-analytic profinite group and let H be a closed subgroup of G. Then the *standard Hausdorff dimension* of H, $\mathrm{hDim}_{\mathrm{Std}}(H)$, is the Hausdorff dimension of H calculated with respect to the natural filtration induced by any given standard open subgroup of G.

We need the following lemma before proving our main theorem.

Lemma 3.3. *Let R be a pro-p ring, and let E be a vector subspace of dimension e of $K^{(d)}$. Then, for every $N \in \mathbb{N}$, the Hausdorff dimension of $(\mathfrak{m}^N)^{(d)} \cap E$ in $R^{(d)}$ with respect to the filtration $\{(\mathfrak{m}^n)^{(d)}\}_{n\in\mathbb{N}}$ is e/d.*

Proof. Let $\mathcal{B} = \{w_1, \dots, w_e\}$ be a basis of E, and let A be the $e \times d$ matrix over K whose rows are the vectors in $\mathcal{B}$. We may assume that A is in reduced echelon form. Set $W = \langle w_1, \dots, w_e \rangle_R$. Let k be such that the product of all denominators of the entries of A belongs to $\mathfrak{m}^k$. One readily checks that

$$\mathfrak{m}^{n+k} W \subseteq (\mathfrak{m}^n)^{(d)} \cap E \subseteq \mathfrak{m}^n W. \tag{5}$$

The required Hausdorff dimension is the limit inferior of the sequence

$$c_n = \frac{\log |(\mathfrak{m}^N)^{(d)} \cap E : (\mathfrak{m}^n)^{(d)} \cap E|}{\log |R^{(d)} : (\mathfrak{m}^n)^{(d)}|},$$

where $n \geq N$. Now, from (5), for $n \geq N + k$, we have

$$|\mathfrak{m}^{N+k} W : \mathfrak{m}^n W| \leq |(\mathfrak{m}^N)^{(d)} \cap E : (\mathfrak{m}^n)^{(d)} \cap E| \leq |\mathfrak{m}^N W : \mathfrak{m}^{n+k} W|.$$

Since W is a free R-module of rank e, we have $\dim_{R/\mathfrak{m}}(\mathfrak{m}^n W/\mathfrak{m}^{n+1} W) = eH(n)$, and consequently

$$\frac{e(H(N+k) + \cdots + H(n-1))}{d(H(1) + \cdots + H(n-1))} \leq c_n \leq \frac{e(H(N) + \cdots + H(n+k-1))}{d(H(1) + \cdots + H(n-1))}.$$

Now, since $H(n) = P(n)$ for large enough n, we have

$$\lim_{n\to\infty} \frac{H(N+k) + \cdots + H(n-1)}{H(1) + \cdots + H(n-1)} = \lim_{n\to\infty} \frac{P(N+k) + \cdots + P(n-1)}{P(1) + \cdots + P(n-1)} = 1,$$

since both sums in the last limit are polynomials in n of degree $\deg P + 1$ and with the same leading coefficient. Arguing similarly with the upper bound for c_n given above, we conclude that $\lim_{n\to\infty} c_n = e/d$, as desired. □

We are now ready to prove our main theorem.

Proof of the Main Theorem. Let $\dim(G) = d$ and $\dim(H) = e$. Since H is an R-analytic submanifold of G, there exist an open subset U of H containing 1, and a chart (V, ϕ) of G such that $U \subseteq V$ and $\phi(U) = E \cap \phi(V)$, for some vector subspace E of $K^{(d)}$ of dimension e.

From the proof of [5, Theorem 13.20], there exist a natural number N and an open subgroup S of G such that $S \subseteq V$ and (S, ϕ) is a standard subgroup of G of level N. Since $U \cap S \subseteq_o U \subseteq_o H$, we have

$$\mathrm{hDim}_{\mathrm{Std}}(H) = \mathrm{hDim}_{\{\mathfrak{m}_\phi^n S\}}(U \cap S),$$

by Lemma 2.4. Since ϕ is an isometry between S and $((\mathfrak{m}^N)^{(d)}, +)$ by Lemma 2.3, we get

$$\mathrm{hDim}_{\{\mathfrak{m}_\phi^n S\}}(U \cap S) = \mathrm{hDim}_{\{(\mathfrak{m}^n)^{(d)}\}}(\phi(U \cap S)).$$

Now since $\phi(U \cap S) = E \cap (\mathfrak{m}^N)^{(d)}$, Lemma 3.3 yields that

$$\mathrm{hDim}_{\{(\mathfrak{m}^n)^{(d)}\}}(\phi(U \cap S)) = \frac{e}{d},$$

which completes the proof. □

Corollary 3.4. *Let G be an R-analytic group of dimension d. Then*

$$\mathrm{Spec}_R(G) \subseteq \left\{0, \frac{1}{d}, \dots, \frac{d-1}{d}, 1\right\}.$$

In particular, the R-analytic spectrum of G is finite and consists of rational numbers.

Acknowledgement. We thank the anonymous referee for helpful comments.

References

[1] J.L. Abercrombie, Subgroups and subrings of profinite rings, *Math. Proc. Cambr. Phil. Soc.* **116** (1994), 209–222.
[2] M. Abért, B. Virág, Dimension and randomness in groups acting on rooted trees, *J. Amer. Math. Soc.* **18** (2005), 157–192.
[3] Y. Barnea, B. Klopsch, Index-subgroups of the Nottingham group, *Adv. Math.* **180** (2003), 187–221.
[4] Y. Barnea, A. Shalev, Hausdorff dimension, pro-p groups, and Kac-Moody algebras, *Trans. Amer. Math. Soc.* **349** (1997), no. 12, 5073–5091.
[5] J. Dixon, M. Du Sautoy, A. Mann, D. Segal, *Analytic pro-p groups*, 2nd ed., Cambridge University Press, Cambridge, 1999.
[6] D. Eisenbud, *Commutative algebra with a view toward algebraic geometry*, Graduate Texts in Mathematics **150**, Springer-Verlag, Berlin-Heidelberg-New York, 2004.
[7] M. Ershov, On subgroups of the Nottingham group of positive Hausdorff dimension, *Comm. Algebra* **35** (2006), 193–206.
[8] K. Falconer, *Fractal geometry: mathematical foundations and applications*, John Wiley and Sons, New York, 1990.
[9] G.A. Fernández-Alcober, A. Zugadi-Reizabal, GGS-groups: order of congruence quotients and Hausdorff dimension, *Trans. Amer. Math. Soc.* **366** (2014), 1993–2017.
[10] J.-P. Serre, *Lie algebras and Lie groups*, Lecture Notes in Mathematics **1500**, Springer-Verlag, Berlin-Heidelberg-New York, 1992.
[11] O. Siegenthaler, Hausdorff dimension of some groups acting on the binary tree, *J. Group Theory* **11** (2008), 555–567.

Department of Mathematics, University of the Basque Country UPV/EHU, 48080 Bilbao, Spain
E-mail address: gustavo.fernandez@ehu.eus

Department of Mathematics, University of Kaiserslautern, P.O. Box 3049, 67655 Kaiserslautern, Germany
E-mail address: gianelli@mathematik.uni-kl.de

Department of Mathematics, University of the Basque Country UPV/EHU, 48080 Bilbao, Spain
E-mail address: jon.gonzalez@ehu.es